



Oligosaccharides Applied to Sugar-conjugates

TCI provides a variety of azido- or amino-linked oligosaccharides and conjugation materials for Sugar-conjugates


Through the research in carbohydrate field, various biological functions of oligosaccharides have been elucidated in life phenomena such as infection, cell-cell recognition, information transmission, fertilization and cell division. The materials containing sugar chains are expected as new functional bioengineering materials which can be applied to medical materials, for example biosensors which specifically bind to toxins or pathogens, composite materials used for surface plasmon resonance (SPR) or crystal oscillator (QCM), or affinity chromatography for the purification of various biopolymers.

Sugar chains having a thiol group, carboxyl group, amino group or azido group at the reducing terminal can be applied to immobilization onto resin beads and gold substrates, introduction of fluorescent groups or biotin, or the use of click chemistry as shown below. Sugar-conjugates are expected as important tools for new bioengineering building materials.

Application 1: Surface Plasmon Resonance (SPR)



Technique to detect interactions or binding capacity between sugars and viruses or proteins.



Biochemical and diagnostic reagents for cancer treatments and missile drugs capable of limited lesion attack.

Application 3: Beads/Resins

Removal of autoantibodies / viruses / toxins and purification of antibody drugs.



Convenient synthesis of sugar-conjugates with proteins and saccharides by click chemistry technique.



Highly sensitive analysis by HPLC, bioimaging and analysis of sugar-protein as well as antigen-antibody interactions.

Oligosaccharides Applied to Sugar-conjugates

TCI products and example of possible assembled sugar-conjugates

Azido-/Amino-linked Oligosaccharides

Sialyl Lewis X-Lactose Ethylazide : Neu5Acα(2-3)Galβ(1-4)[Fucα(1-3)]GlcNAcβ(1-3)Galβ(1-4)Glcβ-ethylazide	[S0922]
Sialyl Lewis X-Lactose Ethylamine : Neu5Acα(2-3)Galβ(1-4)[Fucα(1-3)]GlcNAcβ(1-3)Galβ(1-4)Glcβ-ethylamine	[S0923]
Neu5Acα(2-3)Galβ(1-4)GlcNAc-β-ethylamine	[N0949]
Neu5Acα(2-6)Galβ(1-4)GlcNAc-β-ethylamine	[N0950]
Neu5Acα(2-3)Gal-β-ethylamine	[N0947]
Neu5Acα(2-6)Gal-β-ethylamine	[N0948]
Ganglioside GM_3 (phyto-type): NeuAca(2-3)Gal β (1-4)Glc-ceramide	[G0422]
Disialylnonasaccharide β-ethylazide	[D4217]
$HNK-1\ Ethylazide: Glc A [3S] \beta (1-3) Gal \beta (1-4) Glc NAc \beta (1-3) Gal \beta (1-4) Glc \beta - Ethylazide$	[H1333]
GlcA[3S]β(1-3)Galβ(1-4)GlcNAcβ(1-2)Man β-ethylazide	[G0372]
GlcNAcβ(1-2)Man β-ethylazide	[G0337]
Gb_3 - β -ethylamine : $Gal\alpha(1-4)Gal\beta(1-4)Glc$ - β -ethylamine	[G0402]
$Gb_3\text{-}\beta\text{-}ethylazide: Gal\alpha(1\text{-}4)Gal\beta(1\text{-}4)Glc\text{-}\beta\text{-}ethylazide}$	[G0403]
$Lac DiNAc\ Dimer\ Ethylazide: GalNAc\beta(1-4)GlcNAc\beta(1-3)GalNAc\beta(1-4)GlcNAc-\beta-ethylazide$	[L0237]
GalNAcβ(1-3)GlcNAc β-Ethylazide	[G0373]
A antigen PEG-trifluoroacetamide : GalNAc α (1-3)[Fuc α (1-2)]Gal- β -PEG-trifluoroacetamide	[A2631]
B antigen PEG-trifluoroacetamide : Gal α (1-3)[Fuc α (1-2)]Gal- β -PEG-trifluoroacetamide	[B4172]
N-GlcNAc-Biotin	[G0297]
2-Azidoethyl 2-Acetamido-2-deoxy-β-D-glucopyranoside	[A2172]
2-Azidoethyl 2,3,4,6-Tetra-O-acetyl-β-D-glucopyranoside	[A2377]
2-Azidoethyl 2-Acetamido-2-deoxy-β-D-galactopyranoside	[A2627]


Conjugation Materials



N-Succinimidyl D-Biotinate

5-FITC (isomer I)

5-Hexynoic Acid [H0882]



4,4'-Dithiodibutyric Acid

Sugar-Conjugates



Neu5Aca(2-6)Galb(1-4)GlcNAc having a thiol group

A antigen on beads

Gb₃ FITC-conjugate

Please ask us when you need any sugar chain besides listed products in this leaflet. TCI synthesizes a wide range of sugar chains to meet your request with advanced proprietary technology.

For further information please refer to our website at www.**TCI**chemicals.com

Ordering and Customer Service

TCI AMERICA

Tel : 800-423-8616 / 503-283-1681 : 888-520-1075 / 503-283-1987 E-mail: Sales-US@TCIchemicals.com

TCI EUROPE N.V.

: +32 (0)3 735 07 00 : +32 (0)3 735 07 01 F-mail: Sales-FU@TCIchemicals.com

TCI Deutschland GmbH

Tel : +49 (0)6196 64053-00 : +49 (0)6196 64053-01 Fax E-mail: Sales-DE@TCIchemicals.com

Tokyo Chemical Industry UK Ltd.

: +44 (0)1865 78 45 60 E-mail: Sales-UK@TCIchemicals.com

TCI Chemicals (India) Pvt. Ltd.

E-mail: Sales-IN@TCIchemicals.com

: 1800 425 7889 / 044-2262 0909 Tel : 044-2262 8902 Fax

梯希爱(上海)化成工业发展有限公司 Tel :800-988-0390 / 021-67121386

: 021-6712-1385

E-mail: Sales-CN@TCIchemicals.com

TOKYO CHEMICAL INDUSTRY CO., LTD.

: +81 (0)3-5640-8878

E-mail: globalbusiness@TCIchemicals.com